## JAN 2 0 7005 JAN 2 0 7005 FORM PROMATION DISCLOSURE STATEMENT BY APPLICANT Sheet 1 of 3 APPLICATION NO.: 09/463,874 ATTY. DOCKET NO.: V0179.70000US00 CONFIRMATION NO.: 6909 APPLICANT: Erich Wanker et al. GROUP ART UNIT: 1649 EXAMINER: Olga N. Chernyshev

## U.S. PATENT DOCUMENTS

Examiner's	Cite	U.S. Patent Document		Name of Patentee or Applicant of Cited	Date of Publication or Issue	
Initials #	No.	Number	Kind Code	Document of Cited	of Cited Document MM-DD-YYYY	

## FOREIGN PATENT DOCUMENTS

Examiner's C	Cite	Foreign Patent Document		nent	Name of Patentee or Applicant of Cited	Date of Publication of	Translation	
	Initials "	No.	Office/ Country	Number	Kind Daniman	• •	Cited Document MM-DD-YYYY	(Y/N)
1								

## OTHER ART — NON PATENT LITERATURE DOCUMENTS

Examiner's Initials #	Cite No	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	Translation (Y/N)
		BATES, G.P. et al., Transgenic models of Huntington's disease. Hum Mol Genet. 1997;6(10):1633-7.	
		BECHER, M.W. et al., Intranuclear neuronal inclusions in Huntington's disease and dentatorubral and pallidoluysian atrophy: correlation between the density of inclusions and IT15 CAG triplet repeat length. Neurobiol Dis. 1998 Apr;4(6):387-97.	
		BEYREUTHER, K. et al., Alzheimer's disease. Tangle disentanglement. Nature. 1996 Oct 10;383(6600):476-7.	
		BOOTH, D.R. et al., Instability, unfolding and aggregation of human lysozyme variants underlying amyloid fibrillogenesis. Nature. 1997 Feb 27;385(6619):787-93.	
		BURKE, J.R. et al., Huntingtin and DRPLA proteins selectively interact with the enzyme GAPDH. Nat Med. 1996 Mar;2(3):347-50.	
De		CAPUTO, C.B. et al., Amyloid-like properties of a synthetic peptide corresponding to the carboxy terminus of beta-amyloid protein precursor. Arch Biochem Biophys. 1992 Jan;292(1):199-205.	
		CAUGHEY, B. and CHESEBRO, B. Prion protein and the transmissible spongiform encephalopaties. Trends Cell Biol. 1997; 7: 56-62.	
		DE ROOIJ, K.E. et al., Subcellular localization of the Huntington's disease gene product in cell lines by immunofluorescence and biochemical subcellular fractionation. Hum Mol Genet. 1996 Aug;5(8):1093-9.	
		DIFIGLIA, M. et al., Aggregation of huntingtin in neuronal intranuclear inclusions and dystrophic neurites in brain. Science. 1997 Sep 26;277(5334):1990-3.	·
		DIFIGLIA, M. et al., Huntingtin is a cytoplasmic protein associated with vesicles in human and rat brain neurons. Neuron. 1995 May;14(5):1075-81.	
		DUYAO, M.P. et al., Inactivation of the mouse Huntington's disease gene homolog Hdh. Science. 1995 Jul 21;269(5222):407-10.	
	-	GEORGALIS, Y. et al., Huntingtin aggregation monitored by dynamic light scattering. Proc Natl Acad Sci U S A. 1998 May 26;95(11):6118-21.	
		GLENNER, G.G. Amyloid deposits and amyloidosis. The beta-fibrilloses (first of two parts). N Engl J Med. 1980 Jun 5;302(23):1283-92 and 1333-1343.	
		GOLDBERG, Y.P. et al., Cleavage of huntingtin by apopain, a proapoptotic cysteine protease, is modulated by the polyglutamine tract. Nat Genet. 1996 Aug;13(4):442-9.	
		HDCRG. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. The Huntington's Disease Collaborative Research Group. Cell. 1993 Mar 26;72(6):971-83.	

/6	763		
JAN 2		HOLMBERG, M. et al., Spinocerebellar ataxia type 7 (SCA7): a neurodegenerative disorder with	
1 3514 4	) E		
法	/قير	HOOGEVEEN, A.T. et al., Characterization and localization of the Huntington disease gene	
15 3 × XX	A PAINTS	product, Hum Mol Genet. 1993 Dec;2(12):2069-73.	
		IGARASHI, S. et al., Suppression of aggregate formation and apoptosis by transglutaminase	
		inhibitors in cells expressing truncated DRPLA protein with an expanded polyglutamine stretch. Nat	
		Genet. 1998 Feb;18(2):111-7.	
		IKEDA, H. et al., Expanded polyglutamine in the Machado-Joseph disease protein induces cell	
	<b> </b>	death in vitro and in vivo. Nat Genet. 1996 Jun;13(2):196-202.	
	Ì	JARRETT, J.T. and LANSBURY, P.T. Seeding "one-dimensional crystallization" of amyloid: a	
		pathogenic mechanism in Alzheimer's disease and scrapie? Cell. 1993 Jun 18;73(6):1055-8.	
		KALCHMAN, M.A. et al., HIP1, a human homologue of S. cerevisiae Sla2p, interacts with	
		membrane-associated huntingtin in the brain. Nat Genet. 1997 May;16(1):44-53.	
		KALCHMAN, M.A. et al., Huntingtin is ubiquitinated and interacts with a specific ubiquitin-	
		conjugating enzyme. J Biol Chem. 1996 Aug 9;271(32):19385-94.	
		LI, XJ. et al., A huntingtin-associated protein enriched in brain with implications for pathology.	
		Nature. 1995 Nov 23;378(6555):398-402.	
		LIM, K. et al., Three-dimensional structure of Schistosoma japonicum glutathione S-transferase	
1		fused with a six-amino acid conserved neutralizing epitope of gp41 from HIV. Protein Sci. 1994	
	4	Dec;3(12):2233-44.	
- 1		MATILLA, A. et al., The cerebellar leucine-rich acidic nuclear protein interacts with ataxin-1.	
	ļ	Nature. 1997 Oct 30;389(6654):974-8. Erratum in: Nature 1998 Feb 19;391(6669):818.	
- 1		ONODERA, O. et al., Toxicity of expanded polyglutamine-domain proteins in Escherichia coli.	
		FEBS Lett. 1996 Dec 9;399(1-2):135-9.	
- 1		PAULSON, H.L. et al., Intranuclear inclusions of expanded polyglutamine protein in	
		spinocerebellar ataxia type 3. Neuron. 1997 Aug;19(2):333-44.	
	.1	PERUTZ, M.F. et al., Glutamine repeats as polar zippers: their possible role in inherited neurodegenerative diseases. Proc Natl Acad Sci U S A. 1994 Jun 7;91(12):5355-8.	
610	/	PORTERA-CAILLIAU, C. et al., Evidence for apoptotic cell death in Huntington disease and	
of c		excitotoxic animal models. J Neurosci. 1995 May;15(5 Pt 2):3775-87.	
1		PRUSINER, S.B. et al., Scrapie prions aggregate to form amyloid-like birefringent rods. Cell. 1983	
· 1		Dec;35(2 Pt 1):349-58.	
		ROIZIN, L. et al., Neuronal nuclear-cytoplasmic changer in Huntington's chorea. J. Neurol. Sci.	
1		1983; 61: 37-47.	
		ROOS, R.A.C. et al., Nuclear membrane indentations in Huntington's chorea. J Neurol Sci. 1983	····
		Sep;61(1):37-47.	
		ROSS, C.A. When more is less: pathogenesis of glutamine repeat neurodegenerative diseases.	
1		Neuron. 1995 Sep;15(3):493-6.	
		RUBINSZTEIN, D.C. et al., Phenotypic characterization of individuals with 30-40 CAG repeats in	<del></del>
- 1		the Huntington disease (HD) gene reveals HD cases with 36 repeats and apparently normal elderly	
		individuals with 36-39 repeats. Am J Hum Genet. 1996 Jul;59(1):16-22.	
		SATHASIVAM, K. et al., Identification of an HD patient with a (CAG)180 repeat expansion and	-
		the propagation of highly expanded CAG repeats in lambda phage. Hum Genet. 1997	
		May;99(5):692-5.	
		SCHATZ, P.J. Use of peptide libraries to map the substrate specificity of a peptide-modifying	
- 1		enzyme: a 13 residue consensus peptide specifies biotinylation in Escherichia coli. Biotechnology	
		(N Y). 1993 Oct;11(10):1138-43.	
1		SCHERZINGER, E. et al., Huntingtin-encoded polyglutamine expansions form amyloid-like protein	
		aggregates in vitro and in vivo. Cell. 1997 Aug 8;90(3):549-58.	
		SHARP, A.H. et al., Widespread expression of Huntington's disease gene (IT15) protein product.  Neuron. 1995 May;14(5):1065-74.	
-+-			
- 1		SITTLER, A. et al., Alternative splicing of exon 14 determines nuclear or cytoplasmic localisation of fmr1 protein isoforms. Hum Mol Genet. 1996 Jan;5(1):95-102.	
<del></del>	+	SITTLER, A. et al., SH3GL3 associates with the Huntingtin exon 1 protein and promotes the	
- 1		formation of polygln-containing protein aggregates. Mol Cell. 1998 Oct;2(4):427-36.	
<del>                                     </del>	+	SKINNER, P.J. et al., Ataxin-1 with an expanded glutamine tract alters nuclear matrix-associated	
- 1		structures. Nature. 1997 Oct 30;389(6654):971-4. Erratum in: Nature 1998 Jan 15;391(6664):307.	
<del></del>			

0 42	SMITH, D.B. et al., Single-step purification of polypeptides expressed in Escherichia coli as fusions with glutathione S-transferase. Gene. 1988 Jul 15;67(1):31-40.	
JAN 2 0 3000 W	TELLEZ-NAGEL, I. et al., Studies on brain biopsies of patients with Huntington's chorea. J Neuropathol Exp Neurol. 1974 Apr;33(2):308-32.	
FRARE!	TOWBIN, H. et al., Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. Proc Natl Acad Sci U S A. 1979 Sep;76(9):4350-4.	
or	TROTTIER, Y. et al., Cellular localization of the Huntington's disease protein and discrimination of the normal and mutated form. Nat Genet. 1995 May;10(1):104-10.	
/	VONSATTEL, JP. et al., Neuropathological classification of Huntington's disease. J Neuropathol Exp Neurol. 1985 Nov;44(6):559-77.	
- /	WANKER, E.E. et al., HIP-I: a huntingtin interacting protein isolated by the yeast two-hybrid system. Hum Mol Genet. 1997 Mar;6(3):487-95.	
7		

EXAMINER:	DATE CONSIDERED:
Clliefmur	02/13/06
/	

[NOTE - No copies of U.S. patents, published U.S. patent applications, or pending, unpublished patent applications stored in the USPTO's Image File Wrapper (IFW) system, are included. See 37 CFR §1.98 and 1287OG163. Copies of all other patent(s), publication(s), unpublished, pending U.S. patent applications, or other information listed are provided as required by 37 CFR §1.98 unless 1) such copies were provided in an IDS in an earlier application that complies with 37 CFR §1.98, and 2) the earlier application is relied upon for an earlier filing date under 35 U.S.C. §120.]

<sup>\*</sup> EXAMINER: Initial if reference considered, whether or notication is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

<sup>\*</sup>a copy of this reference is not provided as it was previously cited by or submitted to the office in a prior application, Serial No. \_\_\_, filed \_\_\_, and relied upon for an earlier filing date under 35 U.S.C. 120 (continuation, continuation-in-part, and divisional applications).